



Phylogenetic relationships of xenodermid snakes (Squamata: Serpentes: Xenodermidae), with the description of a new genus

V. Deepak^{1,2,*}, Samuel Lalronunga³, Esther Lalhmingliani³, Abhijit Das⁴, Surya Narayanan⁵, Indraneil Das⁶, David J. Gower²

¹ Senckenberg Dresden, Königsbrücker Landstraße 159, 01109 Dresden, Germany

² Department of Life Sciences, The Natural History Museum, London SW7 5BD, UK

³ Systematics and Toxicology Laboratory, Department of Zoology, Mizoram University, Aizawl 796004, Mizoram, India

⁴ Wildlife Institute of India, Chandrabani, Dehradun 248001, Uttarakhand, India

⁵ Suri Sehgal Centre for Biodiversity and Conservation, Ashoka Trust for Research in Ecology and the Environment, Royal Enclave, Srirampura, Bangalore, Karnataka 560064, India

⁶ Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia

<http://zoobank.org/386BF265-9338-409A-9A44-15B70C953A53>

Corresponding author: V. Deepak (veerappandeepak@gmail.com)

Academic editor Uwe Fritz

Received 1 October 2021

Accepted 3 November 2021

Published 26 November 2021

Citation: Deepak V, Lalronunga S, Lalhmingliani E, Das A, Narayanan S, Das I, Gower DJ (2021) Phylogenetic relationships of xenodermid snakes (Squamata: Serpentes: Xenodermidae), with the description of a new genus. *Vertebrate Zoology* 71 747–762. <https://doi.org/10.3897/vz.71.e75967>

Abstract

Xenodermidae is a generally poorly known lineage of caenophidian snakes found in South, East and Southeast Asia. We report molecular phylogenetic analyses for a multilocus data set comprising all five currently recognised genera and including new mitochondrial and nuclear gene sequence data for the recently described *Stoliczka vanhnuailianai*. Our phylogenetic results provide very strong support for the non-monophyly of *Stoliczka*, as presently constituted, with *S. borneensis* being more closely related to *Xenodermus* than to the Northeast Indian *S. vanhnuailianai*. Based on phylogenetic relationships and morphological distinctiveness, we transfer *Stoliczka borneensis* to a new monotypic genus endemic to Borneo, *Paraxenodermus* **gen. nov.** We also present new morphological data for *P. borneensis*.

Key words

Borneo, endemic, morphology, *Paraxenodermus* gen. nov., phylogeny, taxonomy

Introduction

The caenophidian snake family Xenodermidae Gray, 1849 includes five currently recognised genera, namely *Achalinus* Peters, 1869, *Fimbrios* Smith, 1921, *Parafimbrios* Teynié, David, Lottier, Le, Vidal & Nguyen, 2015, *Xenodermus* Reinhardt, 1836 and *Stoliczka* Jerdon, 1870. *Achalinus* is the most speciose of these genera, with 19

currently recognised species, 10 of which were described in the past five years (Uetz et al. 2021). *Achalinus* spp. are distributed from north of 20° latitude in Vietnam, across south-east China and into central Japan (Fig. 1). *Fimbrios* comprises two species (Smith 1921; Ziegler et al. 2008), distributed in southern and central Laos and Vietnam,

with a record from southern Cambodia (Fig. 1). The two species of the recently described *Parafimbrios* are thus far recorded only from northern Vietnam, Laos and Thailand (Fig. 1). The monotypic *Xenodermus* may be the most widespread xenodermid species, occurring in southernmost Myanmar, Thailand, peninsular Malaysia, Borneo, Sumatra and Java (Fig. 1). The genus *Stoliczka* currently includes three poorly known species with a particularly disjunct distribution, two occurring in Northeast India (*S. khasiensis* Jerdon, 1870 and *S. vanhnuailianai* Lalronunga, Lalmangaiha, Zosangliana, Lalmingliani, Gower, Das & Deepak, 2021) and one in northern and western Borneo (*S. borneensis*) (Das 2021; Stuebing et al. 2014) (Fig. 1). Previously, molecular data were available only for *S. borneensis* (Vidal and Hedges 2002), and few morphological data for the <10 reported specimens of *Stoliczka* (sensu lato) had been published (Lalronunga et al. 2021). In this paper, we report the first molecular data for Northeast Indian *Stoliczka* and new morphological data for *S. borneensis*. We test the monophyly of *Stoliczka*, and describe a new genus for the Bornean species.

Materials and Methods

DNA extraction and amplification

We collected a liver sample from the holotype (and only reported specimen of) *Stoliczka vanhnuailianai*, preserved it in 99% ethanol and stored in -20°C freezer. We extracted genomic DNA using the DNeasy (Qiagen) blood and tissue kit and amplified partial sequences of four mitochondrial (mt) and two nuclear (nu) genes. The mt genes are 16S rRNA (16S, 528 base pairs [bp]), 12S rRNA (12S, 317 bp), cytochrome *b* (*cytb*, 654 bp) and cytochrome oxidase subunit 1 (*col*, 710 bp); and the nu markers are oocyte maturation factor (*cmos*, 449 bp) and neurotrophin-3 (*nt3*, 507 bp). PCR conditions followed previously reported protocols (16S, primers 16Sar-L and 16Sbr-H: Palumbi et al. 1991; 12S, primers 12Sa-L and 12Sb-H: Palumbi et al. 1991; *cytb*, primers GluDG L: Palumbi 1996 and H16064: Burbrink et al. 2000; *col*, primers LCO 1490 (F) and HCO 2198 (R): Folmer et al. 1994; *cmos*, primers S77 and S78: Lawson et al. 2005; *nt3*, primers nt3f and nt3r: Townsend et al. 2008). Sanger sequencing was carried out using the same primers. We assembled contigs from bidirectional sequence chromatograms and edited them in SnapGene Viewer (http://www.snapgene.com/products/snapgene_viewer).

Phylogeny

We aligned the new sequences for *Stoliczka vanhnuailianai* with eight other xenodermids, and an outgroup, the non-xenodermid caenophidian *Acrochordus granulatus*. We checked for stop codons in unexpected regions by translating nucleotide alignments to amino acids for pro-

tein-coding genes (*cytb*, *col*, *cmos*, *nt3*) using MEGA 7 (Kumar et al. 2016). We aligned sequences using ClustalW (Thompson et al. 1994) in MEGA 7 (Kumar et al. 2016) with default settings (alignments online from the Natural History Museum data portal: <https://doi.org/10.5519/gbzyjuli>).

First, we built individual gene trees using Maximum Likelihood (ML). Based on availability of sequence data, we selected one species per xenodermid genus (though included both *S. borneensis* and *S. vanhnuailianai* for *Stoliczka*) and the outgroup. We then aligned and concatenated the six gene sequences into a single dataset (3122 basepairs in length) with ten tips, including the outgroup (Table 1).

We used PartitionFinder2 (Lanfear et al. 2017) to identify the best-fit partition scheme for the concatenated dataset and the best-fit model of sequence evolution for each partition as determined by the Bayesian Information Criterion (BIC), using the default greedy algorithm linked to branch lengths (Lanfear et al. 2012). The best-fit scheme for the concatenated dataset comprises six partitions, by gene and by codon position (Table 2). We performed Maximum Likelihood (ML; Felsenstein 1981) phylogenetic analyses with RAXML GUI Ver. 2.0 (Edler et al. 2021), using the GTRGAMMA model of sequence evolution, which is recommended over GTR+G+I because the 25 rate categories account for potentially invariant sites (Stamatakis 2006). For Bayesian (BI) phylogenetic analyses we used MrBayes 3.2.6 (Ronquist et al. 2012) via the XSEDE portal CIPRES Science Gateway v3.3 (Miller et al. 2010), with default prior settings and with all six partitions assigned their best-fit model as determined by PartitionFinder (Table 2). We set up two separate runs with four Markov chains each, initiated from random trees and allowed to run for one million generations, sampling every 100 generations and discarding the first 25% of trees as “burn-in”. We terminated the analyses when the standard deviation of split frequencies was less than 0.005, and then constructed majority rule consensus trees. We checked for effective sample size (ESS) values using Tracer 1.7 (Rambaut et al. 2014), all parameter values had ESS values >200. We quantified support for internal branches in ML and BI trees using bootstrap (500 replicates) and posterior probability, respectively. We assessed levels of support for relationships incompatible with optimal trees by inspecting bipartition tables of ML bootstrap or BI posterior probability trees using PAUP* 4.0a 169 (X86) (Swofford 2003). We rooted the trees with *Acrochordus granulatus* because it is a non-xenodermid caenophidian snake (Figuroa et al. 2016; Deepak et al. 2018; Zaher et al. 2019).

Molecular dating

We aligned a larger dataset with 68 tips including two scolecophidians (*Gerrhopilus mirus* and *Liotyphlops albirostris*) and representatives of all subfamilies of Alethinophidia, including nine xenodermids (sampling all five currently recognised genera). We aligned this dataset

Table 1. GenBank accession numbers and voucher numbers for the sequences used in this study. Sequences used in the ML and BI concatenated phylogeny are indicated with an asterisk. Accession codes for sequences newly generated in this study are in bold text.

Species	Family	12S	16S	cytb	cmos	mt3	col
<i>Acrochordus granulatus</i> *	Acrochordidae	AF544738	AF544786	AF217841	HM234057	FJ434082	MH273113
<i>Acrochordus javanicus</i>	Acrochordidae	KX694587	AF512745	KX694897	HM234058	KX694991	LC533890
<i>Agkistrodon contortrix</i>	Viperidae (Crotalinae)	AF156587	AF156566	EU483383	—	—	MN135583
<i>Ahaetulla pulverulenta</i>	Colubridae (Ahaetuliinae)	KC347304	KC347339	KC347454	KC347378	—	—
<i>Anilius scytale</i>	Aniliidae	AF544753	FJ755180	U69738	AF544722	FJ434066	—
<i>Anomochilus leonardi</i>	Cylindrophiliidae + Anomochiliidae	AY953430	AY953431	—	—	—	—
<i>Aparallactus capensis</i>	Atractaspididae (Aparallactinae)	FJ404129	AY188045	AY188006	AY187967	—	—
<i>Aplopeltura boa</i>	Parcidae (Pareinae)	AF544761	AF544787	JF827673	JF827696	FJ434085	—
<i>Ashenodipsas laevis</i>	Parcidae (Pareinae)	—	KX660197	KX660469	KX660336	—	—
<i>Azemioops feae</i>	Viperidae (Azemiopinae)	KX694579	AF057234	AY352747	AF544695	KX694977	KP403570
<i>Bitis nasicornis</i>	Viperidae (Viperinae)	DQ305411	AY188048	DQ305457	AY187970	—	MH273549
<i>Boa constrictor</i>	Boidae	AF512744	AB177354	AB177354	AF544676	—	MH140079
<i>Boaedon fuliginosus</i>	Lamprophiidae (Lamprophiinae)	FJ404169	AY188079	AF471060	FJ404270	FJ434094	AY122663
<i>Bothrolycus ater</i>	Lamprophiidae (Lamprophiinae)	FJ404144	AY611859	AY612041	FJ404347	—	MH273562
<i>Bufo depressiceps</i>	Lamprophiidae incertae sedis	FJ404147	AY611860	AY612042	AY611951	—	—
<i>Bufo procterae</i>	Lamprophiidae incertae sedis	FJ404148	AY611818	AY612001	AY611910	—	—
<i>Bungarus fasciatus</i>	Elapidae	EU547135	EU579523	EU579523	AY058924	KX694998	KY769767
<i>Calabaria reinhardtii</i>	Calabariidae	KF576842	Z46494	AY099985	AF544682	—	MH273568
<i>Calamaria pavementata</i>	Colubridae (Calamariinae)	MH445959	KX694624	AF471081	AF471103	KX694999	MK064858
<i>Candoia carinata</i>	Candoiidae	AF544741	EU419850	AY099984	AY099961	FJ434077	—
<i>Cantoria violacea</i>	Homalopsidae	EF395873	KX694627	EF395897	—	KX695001	—
<i>Casarea dussumieri</i>	Bolyeridae	AF544754	AF544827	U69755	AF544731	FJ434069	—
<i>Charina bottae</i>	Charinidae (Charininae)	AF544743	AF544816	AY099986	AY099971	FJ434079	—
<i>Chilabothrus striatus</i>	Boidae	—	—	KC329933	KC329991	DQ465554	—
<i>Contia tenuis</i>	Colubridae (Dipsadinae)	AY577021	AY577030	GU112384	AF471134	—	KU986070
<i>Corallus annulatus</i>	Boidae	JX244286	—	KC750012	KC750007	—	MH140107
<i>Cylindrophis ruffus</i>	Cylindrophiliidae + Anomochiliidae	MK065683	AB179619	AB179619	AF471133	MK064906	—
<i>Daboia russelii</i>	Viperidae (Viperinae)	DQ305413	EU913478	EU913478	AF471156	—	GQ225661
<i>Dityophis vivax</i>	Lamprophiidae	FJ404150	AY188052	AY188013	—	—	KU567322
<i>Epicrates cenchria</i>	Boidae	AF368059	—	HQ399501	KC330008	JX576186	—
<i>Eryx colubrinus</i>	Erycidae	AF544747	AF544819	U69811	AF544716	DQ465569	—
<i>Eryx conicus</i>	Erycidae	GQ225680	AF512743	GQ225658	—	—	—
<i>Eumeces notaeus</i>	Boidae	AF368057	AM236347	HQ399499	HQ399536	—	—
<i>Gerrhopilus mirus</i>	Gerrhopilidae	AM236345	AM236345	AM236345	—	GU902566	AM236345
<i>Grayia ornata</i>	Colubridae (Grayiinae)	AF158434	AY611866	AY612048	AF544684	KX695019	MH274058

Species	Family	12S	16S	cytb	cmos	mt3	col
<i>Holgerharm philippinum</i>	Lamprophiidae (Cyclocorinae)		—	MG458758	MG458766	—	—
<i>Homoroselaps lacteus</i>	Lamprophiidae (Atractaspidinae)	KX694590	AY611809	AY611992	AY611901	KX695021	—
<i>Liasis mackloti</i>	Pythonidae	EF545024	EF545051	U69839	AF544726	FJ434075	—
<i>Liopholidophis sexlineatus</i>	Lamprophiidae (Pseudoxyrhophiinae)	FJ404174	AY188063	DQ979985	AY187985	—	JQ909421
<i>Liotyphlops albostriatus</i>	Anomalepididae	AF366693	AF366762	AF544672	AF544727	—	MH140260
<i>Loxocemus bicolor</i>	Loxocemidae	AF512737	AF544828	AY099993	AY444035	FJ434072	—
<i>Microelaps bicoloratus</i>	Lamprophiidae (Aparallactinae)		—	DQ486349	DQ486173	—	—
<i>Mimophis malfatensis</i>	Lamprophiidae (Psammophiinae)	KX694543	AY188070	DQ486461	AY187992	KX695030	JQ909482
<i>Naja (Afronaja) mossambica</i>	Elapidae	GQ359658	AY611813	AY611996	AY611905	—	—
<i>Naja (Boulengerina) melanoleuca</i>	Elapidae	U96801	AY611812	AY611995	AY611904	—	MH274485
<i>Oxyrhabdium leporinum</i>	Lamprophiidae (Cyclocorinae)	—	—	AF471029	DQ112081	—	—
<i>Oxyuramus scutellatus</i>	Elapidae	EU547100	EU547149	EU547051	EU546916	—	—
<i>Pareas carinatus</i>	Pareidae (Pareinae)	AF544773	AF544802	JF827677	JF827702	FJ434086	—
<i>Prosymna janii</i>	Lamprophiidae (Prosymninae)	FJ404193	FJ404222	FJ404319	FJ404293	—	—
<i>Pseudaspis cana</i>	Lamprophiidae (Pseudaspidinae)	FJ404187	AY611898	AY612080	DQ486167	—	—
<i>Pseudoxenodon karlschmidti</i>	Colubridae (Pseudoxenodontinae)	KX694578	JF697330	AF471080	AF471102	KX695042	MK064781
<i>Python bivittatus</i>	Pythonidae	KF010492	KF010492	JX401131	AF435016	—	KF010492
<i>Rhinophis drummondhayi</i>	Uropeltidae	AY700997	AY701028	AF544673	AF544719	FJ434071	—
<i>Sanzinia madagascariensis</i>	Sanziniidae	EU403571	AY336066	U69866	EU403580	—	MH274606
<i>Tropidophis feicki</i>	Tropidophiidae	AF512733	AF512733	KF811124	KF811110	—	—
<i>Ungaliophis continentalis</i>	Charinidae (Ungaliophiinae)	AF512741	AF544833	U69870	AF544724	FJ434081	—
<i>Xenopeltis unicolor</i>	Xenopeltidae	AF512735	AB179620	AB179620	AF544689	FJ434073	MK064839
<i>Xenophidion schaeferi</i>	Xenophidiidae	—	—	AY574279	MK070320	MK070322	—
<i>Xylophis perroteti</i>	Pareidae (Xylophiinae)	—	MK340908	MN970042	MK344193	—	—
<i>Achalinus rufescens</i> *	Xenodermidae	KX694570	KX694613	KX694895	—	KX694990	—
<i>Achalinus spinalis</i> *	Xenodermidae	MK065581	MK194153	MK201476	—	—	MK064822
<i>Achalinus zugorum</i> *	Xenodermidae	—	MT503100	MT513238	—	—	MT502775
<i>Fimbrios klossi</i> *	Xenodermidae	—	—	KX694894	—	—	KP410745
<i>Parafimbrios lao</i> *	Xenodermidae	—	—	—	—	—	KP410746
<i>Parafimbrios vietnamensis</i> *	Xenodermidae	—	—	—	—	—	MH884515
<i>Stoliczka "borneensis"</i> *	Xenodermidae	AF544779	AF544808	—	AF544721	FJ434083	—
<i>Stoliczka vanhualianai</i> *	Xenodermidae	OL352693	OL352694	OL422473	OL422475	OL422474	OL422476
<i>Xenodermus javanicus</i> *	Xenodermidae	AF544781	AF544810	AF544810	AF544711	—	—

Table 2. Partitions and models of sequence evolution used in the ML and BI phylogenetic analyses for the concatenated dataset. 1st, 2nd and 3rd refer to the codon position.

Partitions	Sites	BI	ML
1	<i>coI</i> ^{1st} , <i>nt3</i> ^{1st}	K80+I	GTR+G
2	<i>cytb</i> ^{2nd} , <i>coI</i> ^{2nd}	HKY+I	GTR+G
3	<i>cytb</i> ^{3rd} , <i>coI</i> ^{3rd}	HKY+G	GTR+G
4	<i>12S</i> , <i>16S</i> , <i>cytb</i> ^{1st}	GTR+G	GTR+G
5	<i>cmos</i> ^{1st} , <i>cmos</i> ^{2nd} , <i>nt3</i> ^{1st} , <i>nt3</i> ^{2nd}	K80+I	GTR+G
6	<i>cmos</i> ^{3rd}	HKY	GTR+G

separately using the same methods outlined above (alignments available at: <https://doi.org/10.5519/gbzyjuli>). We applied seven fossil calibrations (Table 3), largely those recommended by Head (2015) and Head et al. (2016) as recently utilized by Deepak et al. (2021). Additionally, we set the root of the tree at a maximum age of 128 Ma and a minimum age of 123 Ma (i.e., Early Cretaceous, to correspond to the approximate age of the Serpentes root (based on point or mean values from Zheng and Wiens, 2016; Miralles et al. 2018; Burbrink et al. 2020). The best-fitting partition scheme and model(s) of sequence evolution identified using PartitionFinder had ten partitions (Appendix 1). Initially we carried out divergence dating by analysing this dataset and partition scheme with BEAST version 2.5 (Bouckaert et al. 2019) using XSEDE in the CIPRES Science Gateway v3.3 (Miller et al. 2010) under a Yule tree process. We assigned a relaxed log-normal clock for each partition of the concatenated BEAST analysis. We set up two independent runs, each employing the Markov Chain Monte Carlo (MCMC) for 100,000,000 generations, sampling every 5,000 trees. We obtained effective sample size (ESS) values using Tracer 1.7 (Rambaut et al. 2014). ESS values were below 100 for the priors and posteriors employing the best-fit model identified using PartitionFinder. We also repeated the analysis implementing the less-complex HKY model for the partitions but otherwise using the same settings. However, in this second analysis, we recovered ESS values above 200 for all the priors and posteriors for the two independent runs.

Morphology

We provide here morphological and meristic data for two specimens of *Stoliczka borneensis* (BMNH 1946.1.15.58 and UNIMAS 8002) and additional published information on unspecified specimens from Stuebing et al. (2014). Total length, snout-vent length and tail length were measured with thread and a ruler to the nearest 1 mm. Other dimensions were recorded with dial calipers, to the nearest 0.1 mm. Bilateral scale counts separated by a comma are reported in left, right order. Ventral scales were counted following Dowling (1951). Length and width of head scales were measured at the longest and the widest points of the respective scale(s). Eye diameter was measured horizontally.

Museum abbreviations

UNIMAS—Universiti Malaysia Sarawak; NHMUK—Natural History Museum, London (specimen numbers have a BMNH prefix); ZSIK—Zoological Survey of India, Kolkata, and ZRC—Herpetofauna and fish fauna collection at Lee Kong Chian Natural History Museum, Singapore.

Results

Phylogeny

The single-gene ML trees are shown in Fig. 2. Depending on taxon sampling (limited by availability of sequence data), generally *S. borneensis* and *S. vanhnuailianai* show close affinities with *Xenodermus* and with *Achalinus*, respectively. Although ML bootstrap support for many relationships are not strong (<90), support for *Stoliczka* monophyly in the four gene trees for which both species were sampled is negligible, being only 25 for *16S* and 0–0.2 for *12S*, *cmos* and *nt3*. The ML and BI trees derived from the concatenated dataset agree in the set of relationships depicted (Fig. 1), with generally moderate (70–90 ML bootstrap; 0.80–0.90 BI posterior probability) to high support (>90 ML; >0.95 BI). Importantly, there is zero bootstrap or posterior probability support for *Stoliczka* monophyly in these latter trees. Instead, the best-supported relationships that are incompatible with this optimal set of relationships for *Stoliczka* spp. are for *Xenodermus javanicus* being more closely related to *Fimbrios* and *Parafimbrios* (ML bootstrap = 20; BI posterior probability = 0) and for *S. borneensis* being more closely related to *Fimbrios* and *Parafimbrios* (ML bootstrap = 5; BI posterior probability = 9). Thus, we conclude that the available DNA sequence data provide good to strong support for *S. borneensis* being more closely related to *Xenodermus* than to *S. vanhnuailianai*, and for *S. vanhnuailianai* being more closely related to *Achalinus* than to *S. borneensis*, and very strong support for non-monophyly of *Stoliczka*.

Morphology

Previously, extensive data were available for only a single vouchers specimen (the holotype, BMNH 1946.1.15.58) of *Stoliczka borneensis* (Lalronunga et al. 2021). Data for an additional specimen (UNIMAS 8002) are presented in Table 4. This specimen agrees with data presented by Lalronunga et al. (2021) corroborating that *S. vanhnuailianai* resembles the type species of the genus, *S. khasiensis* much more closely than either does *S. borneensis*. Notable differences between the Bornean species and the two Northeast Indian species include presence of 4–6 small scales between the frontals and prefrontals in *S. borneensis* which are absent in the Northeast Indian

Table 3. Parameter values for fossil calibrations used in the BEAST divergence dating analysis. Ages in Ma. All maximum ages soft, except hard maximum for calibration 6.

Calibration	Node calibrations	Offset	Maximum age	Mean	Standard deviation
1	Oldest divergence within crown Alethinophidia	93.9	100.5	1.5	1.25
2	Oldest divergence between non-xenodermid colubroids and their closest living relative (Xenodermidae)	50.5	72.1	6.1	1.25
3	Divergence between Boinae and its sister taxon (Erycinae + Candoiinae)	58	64	1.8	1.25
4	Divergence between <i>Corallus</i> and (<i>Chilabothrus</i> + (<i>Epicrates</i> + <i>Eunectes</i>))	50.2	64	4	1.25
5	Divergence between Viperinae and Crotalinae	20	23.8	1	1.25
6	Divergence between <i>Acrochordus javanicus</i> and (<i>A. ararfu-rae</i> + <i>A. granulatus</i>)	18.1	23.1	1.5	1.25
7	Oldest divergence between <i>Naja (Afronaja)</i> and <i>Naja (Boulengerina)</i>	17	20	1	1.25

Table 4. Morphometric (in mm) and meristic data for *Paraxenodermus borneensis*. Data for the holotype (*) from Lalronunga et al. (2021). Data for unspecified specimens from Stuebing et al. (2014).

Voucher Number	UNIMAS 8002	BMNH 1946.1.15.58*	Unspecified specimens
Sex	male	male	
Snout-vent length	481	541	
Tail length (Ta)	232	248	
Total length (TL)	713	789	
TaL / TL	0.33	0.31	
Horizontal eye diameter	2.7	2.8	
Head length	9.6	17.9	
Head width	8.0	10.8	
Head height	5.2	5.5	
Dorsal scale rows at one head length behind head	31	31	
Dorsal scale rows at midbody	31	32	31–35
Dorsal scale rows at one head length before vent	25	25	
Ventrals	206	208	205–210 (“females only”)
Subcaudals	128	123	117–124
Anal shields	1	1	
Supralabials	10,10	10,-	10 or 11
Supralabials touching eye	0	0	
Infralabials	14,13	14,-	
Infralabials touching anterior genials	1–2	1–3	
Suboculars	3,3	3,3	
Loreals	1,1 (+ 2 very small scales on both sides close to nasals)	1,1 (+ 2 very small scales on both sides close to nasals)	
Preoculars	2,2	3,3	
Supraoculars	3,3	2,2	
Postoculars	4,4	2,4	
Anterior temporals	0,0	0,0	

species; supralabials not contacting the eye in *S. borneensis* versus contacting the eye *S. vanhnuailianai* and *S. khasiensis*; 10 or 11 supralabials versus 8 or 9 supralabials. Although *S. borneensis* is seemingly most closely related to *Xenodermus* (Fig. 2), the two taxa differ markedly in external morphology—for example, *X. javanicus* lacks large scales on the head other than at the snout tip

whereas *S. borneensis* additionally has large parietal and frontal shields. *Xenodermus javanicus* and *S. borneensis* share a derived condition of having more small, irregular head scales than are present in other xenodermids.




Figure 1. Left panel: Multilocus ML phylogeny showing relationships of xenodermid snakes. Numbers at internal branches are: ML bootstrap support / BI posterior probability support. Right panel: map depicting distribution of all currently recognised xenodermid genera. Source: GBIF, Teynié et al. 2015, Ziegler et al. 2008). Additional localities for *Xenodermus javanicus* from Smith, 1943; Tweedie, 1954; Taylor, 1965; David and Vogel, 1996; Wallach et al. 2014).

Systematics

Stoliczka — (Jerdon, 1870)

Stoliczkaia — Boulenger, 1890

Stolickaia — Palacky, 1898

Stolickaia — Palacky, 1898

Estoliczkaia — Briceño-Rossi, 1934

Stoliczkaia — Smith, 1943

Stolickia — Taub, 1967

Stoliczkai — Murthy and Pillai in Majumuria, 1986

Content—*S. khasiensis* (Fig. 3A–B) and *S. vanhnuailianai* (Fig. 3C–D)

Diagnosis. This genus can be diagnosed based on the combination of the following features: (1) maxillary teeth small and subequal, (2) head very distinct from (much wider than) ‘neck’, with large shields on dorsal aspect, (3) posterior one-third of the head and posterior temporal region covered with small scales like those of the anterior end of the body, (4) 3 small scales between parietal and supralabial shields immediately behind eye (5) 8–9 supralabials, (6) nostril in a large concave nasal, (7) body slender and somewhat laterally compressed, (8) ventrals large, and (9) dark dorsum and pale venter meet along a regular straight line ventrolaterally and subcaudals partially or completely darker than venter.

Distribution. This genus is restricted to Northeast India (Fig. 1). *Stoliczka khasiensis* is thus far known only

from Khasi hills, Meghalaya state, India and the recently described *Stoliczka vanhnuailianai* is known only from Mizoram state, India.

Etymology. The genus is named after the Moravian-born Ferdinand Stoliczka (1838–1874). A geologist-natural historian, he was appointed as a palaeontologist with the Geological Survey of India in 1863. Stoliczka collected vertebrates and molluscs from northern India, Andaman and Nicobar Islands, Myanmar and the Malay Peninsula. He served as the official Naturalist with the Second Mission to Yarkand, in central Asia. A biography and a list of published works and reports by Stoliczka can be found in Kolmaš (1982).

Paraxenodermus, gen. nov.

<http://zoobank.org/A08F93FC-187B-48BD-8AFA-A02-EBA98651B>

Type species. *Paraxenodermus borneensis* (Boulenger, 1899).

Type locality. Mount Kinabalu, North Borneo (4,200 ft / 1,280 m); the holotype is deposited in the Natural History Museum, London as BMNH 1946.1.15.58; collected by Richard Hanitsch in March, 1899.

Content—*Paraxenodermus borneensis*




Figure 2. Single-gene ML trees showing inferred phylogenetic relationships of xenodermid snakes, rooted with outgroup *Acrochordus granulatus*. ML bootstrap support is shown at internal branches. Scale bars indicate substitutions per site.

Paraxenodermus borneensis

Figs 3E–F, 4 & 5

Stoliczkaia borneensis Boulenger 1899: 452
Stoliczkaia borneensis — de Rooij 1917: 45
Stoliczkaia borneensis — de Haas 1950: 530
Stoliczkaia borneensis — Haile 1958: 766
Stoliczkaia borneensis — Stuebing: 1991: 329
Stoliczkaia borneensis — Manthey and Grossmann 1997: 394
Stoliczkaia borneensis — Malkmus et al. 2002
Stoliczkaia borneensis — Das 2006a: 9
Stoliczkaia borneensis — Das 2006b: 500–501
Stoliczkaia borneensis — Das 2012: 153
Stoliczkaia borneensis — Das 2018: 151, 169
Stoliczkaia borneensis — Stuebing et al. 2014: 79
Stoliczkaia borneensis — Wallach et al. 2014: 689
Stoliczkaia borneensis — Boundy 2020: 172
Stoliczkaia borneensis — Lalrununga et al. 2021: 569–580

Diagnosis. This genus can be diagnosed based on the combination of the following features: (1) maxillary teeth small and subequal, (2) head very distinct from (much wider than) ‘neck’, with large shields on dorsal aspect,

(3) posterior one-third of the head and posterior temporal region covered with small scales like those of the anterior of the body, (4) numerous small scales between parietal and supralabial shields immediately behind eye, (5) a row of 4–6 small scales between the frontal and prefrontal shields, (6) 10–11 supralabials, (7) nostril in a large concave nasal, (8) body slender and somewhat laterally compressed, (9) ventrals large, and (10) dorsum with numerous dorsolateral and middorsal pale blotches, venter pale with brown patches and subcaudals dark grey.

Comparison to other xenodermid genera. Morphologically *Paraxenodermus borneensis* differs from all other xenodermid snakes by a combination of the following characters: presence of head shields (absent in *Xenodermus javanicus*, other than at snout tip), approximately one-third of the head covered with small scales similar to dorsal scales on the anterior of the body (versus head scales distinct from body scales in *Achalinus*, *Fimbrios* and *Parafimbrios*), head much wider than ‘neck’ (versus head indistinct from neck in *Fimbrios*, *Parafimbrios* and *Achalinus*) and presence of a row of small scales between frontal and prefrontal scales (absent in *Stoliczkaia*).




Figure 3. Line drawings of *Stoliczka khasiensis* (A, B), *Stoliczka vanhuailianai* (C, D) and *Paraxenodermus borneensis* (E, F) based on ZSIK 14945, BNHS 3656 and BMNH 1946.1.15.58 respectively. Genus characteristics are highlighted in different colours: 1) some supralabials in contact with eye in *Stoliczka*, separated by circumorbital scales in *Paraxenodermus*; 2) fewer supra- and infralabials in *Stoliczka* than in *Paraxenodermus*; 3) single prefrontal in *Stoliczka* versus 2–3 in *Paraxenodermus*, 4) fewer scales between parietal and supralabials immediately behind eye in *Stoliczka* than in *Paraxenodermus*, and 5) small row of scales between frontal and prefrontals absent in *Stoliczka*, present in *Paraxenodermus*. Note small scales behind the temporals are indicative rather than precisely accurate. Pale grey coloured areas are bare skin exposed between scales. Illustrations by V. Deepak and Surya Narayanan. Scale bars = 10 mm.

Distribution. The new genus is restricted to the island of Borneo and so far, reported from the Kinabalu Massif (Boulenger 1899) and the contiguous Crocker Range, both in Sabah, in the northeastern part of Borneo (Das 2006a), as well as in the isolated Gunung Murud (Das 2006b), in Sarawak State. https://www.inaturalist.org/observations?taxon_id=28573. Information is not available for the holotype, but all other reported individuals were found late at night, moving slowly on rocky banks of streams at elevations of 950–2,100 m above sea level (Das 2006a).

Variation. The two examined specimens of *Paraxenodermus borneensis*, the holotype BMNH 1946.1.15.58 and UNIMAS 8002, differ slightly in the number of small scales lying between the frontal and prefrontals, being six and four, respectively. We counted six small scales in this position in images of a live individual on the internet (<https://www.inaturalist.org/taxa/28573>). UNIMAS 8002 also differs from BMNH 1946.1.15.58 in having a two more ventrals (208

versus 206) and five additional subcaudals (128 versus 123), and in being smaller (713 mm versus 789 mm total length).

Etymology. The generic name *Paraxenodermus* is composed of the modern Latin generic name *Xenodermus* and the Latin adjective *par* (paris), meaning, among other possibilities, “similar to”.

Discussion

Taken at face value, our phylogenetic results and the distribution of xenodermid genera (Fig. 1) indicate that there are two main radiations within Xenodermidae; one in Northeast India, northern mainland Southeast Asia and Japan (*Stoliczka* + *Achalinus* sensu stricto) and one in eastern mainland Indochina and southeast Sundaland



Figure 4. Holotype of *Paraxenodermus borneensis* (Boulenger, 1899), BMNH 1946.1.15.58. Photographs by Kevin Webb. Scale bar increments in mm.



Figure 5. *Paraxenodermus borneensis* in life (ZRC 2.5731), from Crocker Range, Sabah, in the north-western Borneo. Sequences for this specimen was published in Vidal and Hedges (2002) and used in this study. Photograph by Indraneil Das.




Figure 6. BEAST chronogram showing estimated divergence times for xenodermid snakes inferred from 68 tips for a concatenated mt and nu dataset. Numbers at internal branches indicate mean divergence ages, with blue bars showing 95% highest posterior density intervals. See Appendix 2 for complete dated phylogeny.

(*Fimbrios*, *Parafimbrios*, *Paraxenodermus*, *Xenodermus*). The most parsimonious interpretation is that the most recent common ancestor of these two main xenodermid radiations occurred in mainland Indochina, suggested by our dating analyses to be approximately 66.7–44.6 Ma (Fig. 6). However, this would be better tested in future by undertaking probabilistic biogeographic analyses of a more broadly taxonomically sampled tree.

Establishment of a new genus for *S. borneensis* and a new understanding of phylogenetic relationships removes the exceptional geographic disjunction presented by the previous concept of *Stoliczka*. These results also strengthen evidence for endemic radiations within both Borneo (e.g. Blackburn et al. 2010; Wood et al. 2012; Hertwig et al. 2013; Fritz et al. 2014) and Northeast India (e.g., Pawar et al. 2007; Kamei et al. 2012).

Acknowledgements

We thank K. Lalhmangaiha and Isaac Zosangliana for their support in the field. Specimen of *S. vanhnuailianai* was collected under the research and collection permission (A.38011/5/2011-CWLW/338) issued by the Department of Environment, Forest and Climate Change, Government of Mizoram. AD's research is supported by SERB-DST (CRG/2018/000790) and Director, Wildlife Institute of India. Special thanks are due to Mal-sawmdawngliana for research assistance. Sabah field work by I. Das was supported by research grant UNIMAS 192/99(4), under research permit from Sabah Parks TS/PTD/5/5 Jld.14(76). UNIMAS 8002 from Sarawak was collected under research grant IRPA 08-02-09-10007-EA0001 by the Ministry of Science, Technology and Environment, Government of Malaysia, under research permit from the Sarawak Biodiversity Centre, SBC-RP-0070-ID. VD's contribution was supported in part by a Humboldt Fellowship. VD thanks Uwe Fritz for his support. SN thanks Aravind N.A for his support at ATREE. Kevin Webb (NHMUK, London) is thanked for expert photography of the holotype of *Stoliczka borneensis*. We thank Pratyush P. Mohapatra ZSI, Jabalpur for sharing photographs of the *Stoliczka khasiensis* specimen stored in ZSI, Kolkata. We thank Lee Grismer, Luru Quang Vinh and an anonymous reviewer for their comments and suggestions which improved this manuscript.

References

- Blackburn DC, Bickford DP, Diesmos AC, Iskandar DT, Brown RM (2010) An ancient origin for the enigmatic flat-headed frogs (Bombinatoridae: Barbourula) from the islands of Southeast Asia. *PLoS One* 5(8): e12090. <https://doi.org/10.1371/journal.pone.0012090>
- Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled J, Jones G, Kühnert D, de Maio N, Matschiner M (2019) BEAST 2.5: An advanced software platform for bayesian evolutionary analysis. *PLoS Computational Biology* 15(4): e1006650. <https://doi.org/10.1371/journal.pcbi.1006650>
- Boulenger GA (1899) Description of three new reptiles and a new batrachian from Mt. Kina Balu, North Borneo. *Annals and Magazine of Natural History Series 7*, 4(24): 451–454. <https://doi.org/10.1080/00222939908678228>
- Boulenger GA (1890) The fauna of British India, including Ceylon and Burma. Reptilia and Batrachia. Taylor and Francis, London, xviii + 541 pp. 142 Figs
- Boundy J (2020) Snakes of the World: a supplement. CRC Press. 273 pp.
- Briceño-Rossi AL (1934) El problema del ofidismo en Venezuela. *Boletín del Ministerio de Salubridad y de Agricultura y Cria*, Caracas 2(14), 1079–1177 & 2(15), 15–103, illust.
- Burbrink FT, Lawson R, Slowinski JB (2000) Mitochondrial DNA phylogeography of the polytypic north American rat snake (*Elaphe obsoleta*): A critique of the subspecies concept. *Evolution* 54(6): 2107–2118. [https://doi.org/10.1554/00143820\(2000\)054\[2107:MD-POTP\]2.0.CO;2](https://doi.org/10.1554/00143820(2000)054[2107:MD-POTP]2.0.CO;2)
- Burbrink FT, Grazziotin FG, Pyron RA, Cundall D, Donnellan S, Irish F, Keogh JS, Kraus F, Murphy RW, Noonan B, Raxworthy CJ, Ruane S, Lemmon AR, Lemmon EM, Zaher H (2020) Interrogating genomic-scale data for Squamata (lizards, snakes, and amphisbaenians) shows no support for key traditional morphological relationships. *Systematic Biology* 69: 502–520. <https://doi.org/10.1093/sysbio/syz062>
- Das I (2006a) Crocker Range National Park, Sabah, as a refuge for Borneo's montane herpetofauna. *Amphibian and Reptile Conservation* 4(1): 3–11. <https://doi.org/10.1514/journal.arc.0040015>
- Das I (2006b) Geographic Distribution. *Stoliczka borneensis* (Stoliczka's Water Snake). *Herpetological Review* 37(4): 500–501.
- Das I (2021) A Naturalist's Guide to the Snakes of South-East Asia: Malaysia, Singapore, Thailand, Myanmar, Borneo, Sumatra, Java and Bali. Third edition. John Beaufoy Publishing. Oxford, 176 pp.
- David P, Vogel G (1996) The snakes of Sumatra. Chimaira, Frankfurt am Main, 260 pp.
- Deepak V, Ruane S, Gower DJ (2018) A new subfamily of fossorial colubroid snakes from the Western Ghats of peninsular India. *Journal of Natural History* 52(45–46): 2919–2934. <https://doi.org/10.1080/00222933.2018.1557756>
- Deepak V, Maddock ST, Williams R, Nagy ZT, Conradie W, Rocha S, Harris DJ, Perera A, Gvoždík V, Doherty-Bone TM, Kamei RG, Menegon M, Labisko J, Morel C, Cooper N, Day JJ, Gower DJ (2021) Molecular phylogenetics of sub-Saharan African natricine snakes, and the biogeographic origins of the Seychelles endemic *Lycognathophis seychellensis*. *Molecular Phylogenetics and Evolution* 161: 107152. <https://doi.org/10.1016/j.ympev.2021.107152>
- de Rooij N (1917) The Reptiles of the Indo-Australian Archipelago. II. Ophidia. Leiden (E. J. Brill), xiv + 334 pp. <https://doi.org/10.5962/bhl.title.5069>
- Dowling HG (1951) A proposed standard system of counting ventrals in snakes. *British Journal of Herpetology* 1(5): 97–99.
- Edler D, Klein J, Antonelli A, Silvestro D (2021) raxmlGUI 2.0: A graphical interface and toolkit for phylogenetic analyses using RAXML. *Methods in Ecology and Evolution* 12: 373–377. <https://doi.org/10.1111/2041-210X.13512>
- Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *Journal of Molecular Evolution* 17: 368–376. <https://doi.org/10.1007/BF01734359>
- Fritz U, Gemel R, Kehlmaier C, Vamberger M, Präsag P (2014) Phylogeography of the Asian softshell turtle *Amyda cartilaginea* (Boddaert, 1770): evidence for a species complex. *Vertebrate Zoology* 64(2): 229–243.
- Figueroa A, McKelvy AD, Grismer LL, Bell CD, Lailvaux SP (2016) A species-level phylogeny of extant snakes with description of a new colubrid subfamily and genus. *PLoS One* 11(9): 1–31. <https://doi.org/10.1371/journal.pone.0161070>
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome C oxidase subunit

- I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299.
- Hass CPJ, De (1950) Checklist of the snakes of Indo-Australian Archipelago. *Treubia* 20: 511–625
- Haile NS (1958) The snakes of Borneo, with a key to the species. *Sarawak Museum Journal* 8: 743–771.
- Head JJ (2015) Fossil calibration dates for molecular phylogenetic analysis of snakes 1: Serpentes, Alethinophidia, Boidae, Pythonidae. *Palaeontologia Electronica* 18: 1–17. <https://doi.org/10.26879/487>
- Head JJ, Mahlow K, Mueller J (2016) Fossil calibration dates for molecular phylogenetic analysis of snakes 2: Caenophidia, Colubroidea, Elapoidea, Colubridae. *Palaeontologia Electronica* 19: 1–21. <https://doi.org/10.26879/625>
- Hertwig ST, Schweizer M, Das I, Haas A (2013) Diversification in a biodiversity hotspot—The evolution of Southeast Asian rhacophorid tree frogs on Borneo (Amphibia: Anura: Rhacophoridae). *Molecular Phylogenetics and Evolution* 68(3): 567–581. <https://doi.org/10.1016/j.ympev.2013.04.001>
- Jerdon TC (1870) Notes on Indian herpetology. *Proceedings of the Asiatic Society of Bengal* 1870: 66–85.
- Kamei RG, Mauro DS, Gower DJ, Van Bocxlaer I, Sherratt E, Thomas A, Babu S, Bossuyt F, Wilkinson M, Biju SD (2012) Discovery of a new family of amphibians from northeast India with ancient links to Africa. *Proceedings of the Royal Society B: Biological Sciences* 279(1737): 2396–2401. <https://doi.org/10.1098/rspb.2012.0150>
- Kolmaš J (1982) Ferdinand Stoliczka (1838–1874): the life and work of the Czech explorer in India and High Asia. *Wiener Studien zur Tibetologie und Buddhismuskunde. Universität Wien, Vienna*. Vol. 9. xi + 58 pp.
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33: 1870–1874. <https://doi.org/10.1093/molbev/msy096>
- Lalronunga S, Lalmangaiha K, Zosangliana I, Lalmingliani E, Gower DJ, Das A, Deepak V (2021) A new species of *Stoliczka* Jerdon, 1870 (Serpentes: Xenodermidae) from Mizoram, India. *Zootaxa* 4996(3): 569–580. <https://doi.org/10.11646/zootaxa.4996.3.9>
- Lanfear R, Calcott B, Ho SY, Guindon S (2012) PartitionFinder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* 29: 1695–1701. <https://doi.org/10.1093/molbev/mss020>
- Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2017) PartitionFinder 2: New methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution* 34: 772–773. <https://doi.org/10.1093/molbev/msw260>
- Lawson R, Slowinski JB, Crother BI, Burbrink FT (2005) Phylogeny of the Colubroidea (Serpentes): New evidence from mitochondrial and nuclear genes. *Molecular Phylogenetics and Evolution* 37(2): 581–601. <https://doi.org/10.1016/j.ympev.2005.07.016>
- Malkmus R, Manthey U, Vogel G, Hoffmann P, Kosuch J (2002) Amphibians and reptiles of Mount Kinabalu (North Borneo). A.R.G. Ganther Verlag, Rugell, 404 pp.
- Manthey U, Grossmann W (1997) Amphibien & Reptilien Südasiens. *Natur und Tier Verlag (Münster)*, 512 pp.
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *Proceedings of the Gateway Computing Environments Workshop (GCE)*, New Orleans, Louisiana, 14 November 2010, 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Miralles A, Marin J, Markus D, Herrel A, Hedges SB, Vidal N (2018) Molecular evidence for the paraphyly of Scolecophidia and its evolutionary implications. *Journal of Evolutionary Biology* 31(12): 1782–1793. <https://doi.org/10.5061/dryad.4p2781m>
- Murthy TSN, Pillai RS (1986) Snakes. In: Majumuria, T.C. (Ed.), *Wild-life wealth of India (resources & management)*. Tecpress Service, Bangkok, 224–244, illust.
- Palacky J (1898) La distribution des ophidiens sur le globe. *Mémoires de la Société Zoologique de France* 11: 88–125.
- Palumbi SR, Martin AP, Romano SL, McMillan WO, Stice L, Grabowski G (1991) *The Simple Fool's Guide to PCR ver. 2*. University of Hawaii, Honolulu: 25–28.
- Palumbi SR (1996) The polymerase chain reaction. In: Hillis DM, Moritz C, Mable BK (Eds) *Molecular systematics*. 2nd ed. Massachusetts: Sinauer Associates, 205–247.
- Pawar S, Koo MS, Kelley C, Ahmed MF, Chaudhuri S, Sarkar S (2007) Conservation assessment and prioritization of areas in Northeast India: Priorities for amphibians and reptiles. *Biological Conservation* 136(3): 346–361. <https://doi.org/10.1016/j.biocon.2006.12.012>
- Peters WCH (1869) Über neue Gattungen und neue oder weniger bekannte Arten von Amphibien (*Eremias*, *Dicrodon*, *Euprepes*, *Lygosoma*, *Typhlops*, *Eryx*, *Rhynchonyx*, *Elapomorphus*, *Achalinus*, *Coronella*, *Dromicus*, *Xenopholis*, *Anoplodipsas*, *Spilotes*, *Tropidonotus*). *Monatsberichte der königlich Akademie der Wissenschaften zu Berlin* 1869(5): 432–445.
- Rambaut A, Suchard MA, Xie D, Drummond AJ (2014) Tracer. Version 1.6. Available from: <http://beast.bio.ed.ac.uk/Tracer> (accessed 27 July 2014).
- Reinhardt JT (1836) Afhandling om *Xenodermus javanicus*. Oversigt over det Kongelige Danske videnskabernes selskabs forhandling. Kjobenhavn 3: 6–7.
- Ronquist F, Teslenko M, van der Mark P, Ayres D, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Smith MA (1921) New or little-known reptiles and batrachians from southern Annam (Indochina). *Proceedings of the Zoological Society of London* 1921: 423–440.
- Smith MA (1943) The fauna of British India, Ceylon and Burma, including the whole of the Indo-Chinese subregion. *Reptilia and Amphibia*. Vol. III, Serpentes. Taylor & Francis, London, xii + 583 pp.
- Stamatakis A (2006) RAXML-VI-HP: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22(21): 2688–2690. <https://doi.org/10.1093/bioinformatics/btl446>
- Stuebing RB (1991) A checklist of the snakes of Borneo. *Raffles Bulletin of zoology* 39(2): 323–362
- Stuebing RB, Inger RF, Tan FL (2014) *Field guide to the snakes of Borneo*. Second Edition. Natural History Publications (Borneo). 310 pp.
- Swofford DL (2003) PAUP: phylogenetic analysis using parsimony, version 4.0a 165. Cary (NC): SAS Institute.
- Taub AM (1965) Comparative histological studies on the parotid gland of colubrid snakes. Doctoral Dissertation, State University of New York, Buffalo, 239 pp., 3 Figs, 8 pls.
- Taylor EH (1965) The serpents of Thailand and adjacent waters. *University of Kansas Science Bulletin* 45(9): 609–1096.

- Teynié A, David P, Lottier A, Le MD, Vidal N, Nguyen TQ (2015) A new genus and species of xenodermatid snake (Squamata: Caenophidia: Xenodermatidae) from northern Lao People's Democratic Republic. *Zootaxa* 3926(4): 523–540. <https://doi.org/10.11646/zootaxa.3926.4.4>
- Thompson JD, Higgins DG, Gibson TJ (1994) ClustalW: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673–4680. <https://doi.org/10.1093/nar/22.22.4673>
- Townsend TM, Alegre RE, Kelley ST, Wiens JJ, Reeder TW (2008) Rapid development of multiple nuclear loci for phylogenetic analysis using genomic resources: an example from squamate reptiles. *Molecular Phylogenetics and Evolution* 47(1): 129–142. <https://doi.org/10.1016/j.ympev.2008.01.008>
- Tweedie MWF (1954) *The Snakes of Malaya*, Government Printing Office, Singapore.
- Uetz P, Freed P, Hošek J (2021) *The Reptile Database*. Available from: <http://reptile-database.reptarium.cz> (accessed 1 March 2021)
- Vidal N, Hedges SB (2002) Higher-level relationships of caenophidian snakes inferred from four nuclear and mitochondrial genes. *Comptes Rendus Biologies* 325(9): 987–995. [https://doi.org/10.1016/S1631-0691\(02\)01509-3](https://doi.org/10.1016/S1631-0691(02)01509-3)
- Wallach, V, Williams KL, Boundy J (2014) *Snakes of the World: A Catalogue of Living and Extinct Species*. Taylor and Francis, CRC Press, 1237 pp.
- Wood PL Jr, Heinicke MP, Jackman TR, Bauer AM (2012) Phylogeny of bent-toed geckos (*Cyrtodactylus*) reveals a west to east pattern of diversification. *Molecular Phylogenetics and Evolution* 65(3): 992–1003. <https://doi.org/10.1016/j.ympev.2012.08.025>
- Zaher H, Murphy RW, Arredondo JC, Graboski R, Machado-Filho PR, Mahlow K, Montingelli GG, Zhang YP (2019) Large-scale molecular phylogeny, morphology, divergence-time estimation and the fossil record of advanced caenophidian snakes (Squamata: Serpentes). *PLoS ONE* 14(5): e0216148. <https://doi.org/10.1371/journal.pone.0216148>
- Zheng Y, Wiens JJ (2016) Combining phylogenomic and supermatrix approaches, and a time-calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes and 4162 species. *Molecular Phylogenetics and Evolution* 94: 537–547. <https://doi.org/10.1016/j.ympev.2015.10.009>
- Ziegler T, David P, Miralles A, Van Kien D, Truong NQ (2008) A new species of the snake genus *Fimbrios* from Phong Nha-Ke Bang National Park, Truong Son, central Vietnam (Squamata: Xenodermatidae). *Zootaxa* 1729(1): 37–48. <https://doi.org/10.11646/zootaxa.1729.1.4>

Appendix 1

Partitions and models of sequence evolution used in the BEAST analyses for the 68 tips dataset. 1st, 2nd and 3rd refer to the codon position.


Partitions	Sites	model
1	<i>12S</i>	GTR+G
2	<i>16S</i> , <i>cytb</i> ^{1st}	GTR+I+G
3	<i>col</i> ^{1st} , <i>cytb</i> ^{2nd}	SYM+I+G
4	<i>cytb</i> ^{3rd}	GTR+I+G
5	<i>col</i> ^{2nd}	HKY+G
6	<i>col</i> ^{3rd}	GTR+I+G
7	<i>cmos</i> ^{1st} , <i>cmos</i> ^{2nd}	K80+G
8	<i>cmos</i> ^{3rd}	HKY+G
9	<i>nt3</i> ^{1st} , <i>nt3</i> ^{2nd}	SYM+G
10	<i>nt3</i> ^{3rd}	HKY+G

Appendix 2

Specimens examined and/or photographed

Achalinus meridianus (holotype) BMNH 1946.1.12.31
Achalinus formosanus (holotype) BMNH 1946.1.7.78
Fimbrios klossi (syntype) BMNH 1946.1.15.87
Xenodermus javanicus (holotype) BMNH 1946.1.15.90
Stoliczka khasiensis (holotype) BMNH 1946.1.15.67
Stoliczka khasiensis ZSIK 14945
Stoliczka borneensis (holotype) BMNH 1946.1.15.58
Stoliczka borneensis UNIMAS 8002, ZRC 2.5731
Stoliczka vanhuailianai (holotype) BNHS 3656

Appendix 3



BEAST chronogram generated using concatenated-gene for representatives of all families and subfamilies of alethinophidian snakes. Error bars and the numbers at internal branches indicate 95% highest posterior densities for node ages.

Appendix 4



Paraxenodermus borneensis (UNIMAS 8002) from near Samling Camp at Ravenscourt, Lawas, Sarawak Malaysia. Photographs by Indraneil Das.